

3. (a) What is space group? (2)
- (b) Derive the dispersion relation for a linear diatomic lattice. Sketch the dispersion relation indicating the two branches clearly in the graph. (4+2)
- (c) What are meant by normal and Umklapp processes? (2)
4. (a) Prove the equivalence between vibrational mode in a solid and a harmonic oscillator. (6)
- (b) What is geometrical structure factor? Show that the factor vanishes unless the number h , k and l are all even or all odd for f.c.c. lattice. (4)

.....

(Internal Assessment – 10)

2016
M.Sc.
2nd Semester Examination
PHYSICS

PAPER – PGS-202 (Gr. – A + B)

Full Marks : 50

Time : 2 Hours

*The figures in the right hand margin indicate full marks.
 Candidates are required to give their answers in their own words
 as far as practicable.*

(Gr. A – Quantum Mechanics I)
Answer Q1 and any one from Q2 and Q3.

1. Answer any five bits: 5 X 2 = 10


(i) A particle of mass m is confined in a two-dimensional square well potential of dimension a . The potential $V(x,y)$ is given by

$$V(x,y) = 0 \text{ for } -a < x < a \text{ and } -a < y < a;$$

$$= \infty \text{ elsewhere.}$$

Find out the energy of the first excited state for this particle.

(ii) A particle is constrained to move in a truncated harmonic potential well ($x > 0$) as shown in the figure. Find out the energy of the first excited state of this particle with explanation.



(Turn Over)

(iii) The wavefunction of a particle is given by $\psi = \frac{\varphi_0}{\sqrt{2}} + \frac{i\varphi_1}{\sqrt{2}}$ where φ_0 and φ_1 are the normalized eigenfunctions with energies E_0 and E_1 corresponding to the ground state and the first excited state, respectively. Calculate the expectation value of the Hamiltonian in the state ψ .

(iv) Find the value of commutator $[L_z, \cos\varphi]$ where φ is the azimuthal angle and $\cos\varphi$ is the operator.

(v) Show that the norm of the state vector evolving from the Schrodinger Picture remains invariant with respect to time.

(vi) Deduce the parity of spherical harmonics $Y_{lm}(\theta, \varphi)$.

(vii) For what values of the constant c will be the function $f(x) = A e^{-\alpha x}$ be an eigenfunction of the operator $\hat{Q} = \frac{d^2}{dx^2} + \frac{2}{x} \frac{d}{dx} + \frac{c}{x}$?

(viii) If \hat{A} and \hat{B} are Hermitian operators, prove that $(\hat{A}\hat{B} + \hat{B}\hat{A})$ is Hermitian but $(\hat{A}\hat{B} - \hat{B}\hat{A})$ is non-Hermitian.

2. (a) Considering the operator $a = \frac{i}{\sqrt{2m\hbar\omega}}(\hat{p} - im\hat{x})$ and its adjoint prove that $H|n\rangle = (n+1/2)\hbar\omega|n\rangle$ for a linear harmonic oscillator where the symbols have their usual meanings. (4)

(b) Sketch the ground state wavefunction of hydrogen atom and hence, obtain the expectation value of r^2 in this state. (3)

(c) Discuss the equations of motion for both the wavefunction and operator in interaction picture.

3. (a) The wavefunction for a particle is represented as $\psi(\vec{r}, t) = \sum_n a_n(t) u_n(\vec{r})$ where $H u_n(\vec{r}) = \epsilon_n u_n(\vec{r})$. Show that $\langle \psi | H | \psi \rangle = \sum_n |a_n(t)|^2 \epsilon_n$. (3)

...Continued

(b) A three level quantum system has energy eigenvalues 0, 1, 2 MeV. If the probabilities for the system, at time t , to be in the first eigenstates are 49% and 36% respectively, write down the wavefunction $\psi(\vec{r}, t)$ for the system.

(2)

(c) Prove that the ground-state wavefunction of a linear harmonic oscillator is Gaussian. (2)

(d) A particle is confined to a box of length L with walls $x = 0$ and $x = L$. The particle is described by a wavefunction $\psi = N \sin(3\pi x/2L) \cos(\pi x/2L)$. Find the energy of the particle. (3)

(Gr. B – Solid State Physics I)

Answer Q1 and Q2 and any one from Q3 and Q4.

1. Answer any two bits: 2 X 2 = 4

(i) Show that five-fold rotational axis does not exist in a lattice system.

(ii) Find the packing fraction of the hcp structure.

(iii) The primitive lattice translation vectors of hexagonal space lattice may be taken as, $\vec{a}_1 = \left(\frac{\sqrt{3}}{2}a\right)\hat{i} + \left(\frac{1}{2}a\right)\hat{j}$; $\vec{a}_2 = -\left(\frac{\sqrt{3}}{2}a\right)\hat{i} + \left(\frac{1}{2}a\right)\hat{j}$; $\vec{a}_3 = c\hat{k}$.

Show that the lattice is its own reciprocal, but with a rotation of axis.

2. Answer any two bits: 3 X 2 = 6

(i) What is Brillouin Zone? How can it be constructed using Bragg's diffraction condition?

(ii) Find the dispersion relation for mono-atomic lattice.

(iii) Find out the structure factor for the basis of diamond and prove that if all indices are even, the structure factor of the basis vanishes unless $h + k + l = 4n$, where n is an integer.

(Turn Over)